arXiv:1407.2829v1 [math.CO] 10 Jul 2014

Sketch of a Proof of an Intriguing Conjecture of Karola Mészáros and Alejandro Morales Regarding the Volume of the D_n Analog of the Chan-Robbins-Yuen Polytope (Or: The Morris-Selberg Constant Term Identity Strikes Again!)

By Doron ZEILBERGER

To Dick Askey (b. June 4, 1933): from a 4^3-year-old to a 3^4-year-old, and thanks for preaching the importance of Constant Term Identities!

Disclaimer: Some of the steps below (in particular, the "change of variable" in contour-integrals) require 'rigorous' justification, that I am sure could be easily supplied by a skilled analyst.

Recall that for any rational function $f(z)$ of a variable z, $CT_z f(z)$ is the coeff. of z^0 in the formal Laurent expansion of $f(z)$ (that always exists!).

Karola Mészáros and Alejandro Morales have recently made the following intriguing conjecture.

Conjecture ([MeMora], Conj. 7.12, also presented by Morales[Mora] at Stanley@70)

$$CT_{x_n} CT_{x_{n-1}} \dots CT_{x_1} \prod_{i=1}^{n} x_i^{-1}(1-x_i)^{-2} \prod_{1\le i<j\le n} (x_j-x_i)^{-1}(1-x_j-x_i)^{-1} \quad = \quad 2^{n^2} \prod_{k=1}^{n} Cat(k) \quad ,$$

where $Cat(k)$ are the ubiquitous Catalan numbers $(2k)!/(k!(k+1)!)$ (that Igor Pak believes are better than primes for searching for ETI!)

The present conjecture is a D_n-analog of a conjecture made in [CRY], and proved in [Z], using the versatile Morris-Selberg Constant Term Identity ([Morr], restated in [Z]):

$$CT_{x_n} \dots CT_{x_1} \prod_{i=1}^{n}(1-x_i)^{-a} \prod_{i=1}^{n} x_i^{-b} \prod_{1\le i<j\le n} (x_j-x_i)^{-2c} = S_n(a,b,c) \quad , \qquad (Chip)$$

where

$$S_n(a,b,c) := \frac{1}{n!} \prod_{j=0}^{n-1} \frac{\Gamma(a+b+(n-1+j)c)\Gamma(c)}{\Gamma(a+jc)\Gamma(c+jc)\Gamma(b+jc+1)} \quad .$$

By using Cauchy's theorem, this is equivalent to

$$(\frac{1}{2\pi i})^n \int_C \prod_{i=1}^{n}(1-x_i)^{-a} \prod_{i=1}^{n} x_i^{-b-1} \prod_{1\le i<j\le n} (x_j-x_i)^{-2c} \prod_{i=1}^{n} dx_i \; = \; S_n(a,b,c) \quad , \qquad (Atle)$$

where *now* a, b, c can be *any* real numbers (with obvious conditions to ensure convergence) and C is *any* multi-contour in n-dimensional complex space, that is far enough from the origin.

The Mészáros-Morales conjecture is the special case $a = 2$, $c = \frac{1}{2}$ of the following fact.

Fact : Let a be a positive integer, and c a positive half-integer, then

$$CT_{x_n}\, CT_{x_{n-1}}\, \ldots\, CT_{x_1}\, \prod_{i=1}^{n} x_i^{-(a-1)}(1-x_i)^{-a} \prod_{1\le i<j\le n} (x_j-x_i)^{-2c}(1-x_j-x_i)^{-2c}$$

$$= 2^{2cn(n-1)+2(a-1)n}\, \cdot\, S_n(a\,, -\frac{1}{2}\,, c) \quad .$$

Proof: Converting the iterated constant-terms to a multi-contour-integral, we have to evaluate

$$(\frac{1}{2\pi i})^n \int_C \prod_{i=1}^{n} x_i^{-a}(1-x_i)^{-a} \prod_{1\le i<j\le n} (x_j-x_i)^{-2c}(1-x_j-x_i)^{-2c} \prod_{i=1}^{n} d\,x_i \quad .$$

Now make the *change of variables*:

$$x_i = \frac{1-z_i}{2} \quad , \quad (1\le i\le n) \quad ,$$

getting that our multi-integral equals

$$(\frac{1}{2\pi i})^n\, \cdot\, 2^{2an} \cdot 2^{2cn(n-1)} (-\frac{1}{2})^n \int_{C'} \prod_{i=1}^{n} (1-z_i)^{-a}(1+z_i)^{-a} \prod_{1\le i<j\le n} (z_i-z_j)^{-2c}(z_i+z_j)^{-2c} \prod_{i=1}^{n} d\,z_i$$

$$= (\frac{1}{2\pi i})^n\, \cdot\, (-1)^n 2^{2an+2cn(n-1)-n} \int_{C'} \prod_{i=1}^{n} (1-z_i^2)^{-a} \prod_{1\le i<j\le n} (z_i^2-z_j^2)^{-2c} \prod_{i=1}^{n} d\,z_i \quad ,$$

where C' is some other multi-contour. Now is time for *yet another* **change of variable**

$$w_i = z_i^2 \quad , \quad (1\le i\le n) \quad ,$$

and we have

$$\prod_{i=1}^{n} d\,z_i = (\frac{1}{2})^n \prod_{i=1}^{n} \frac{d\,w_i}{w_i^{1/2}} \quad ,$$

getting that our multi-integral is

$$(-1)^{n+n(n-1)c} (\frac{1}{2\pi i})^n\, 2^{2an+2cn(n-1)-2n} \int_{C''} \prod_{i=1}^{n} (1-w_i)^{-a} \prod_{i=1}^{n} w_i^{-1/2} \prod_{1\le i<j\le n} (w_j-w_i)^{-2c} \prod_{i=1}^{n} d\,w_i \quad ,$$

for yet another multi-contour, C'', and thanks to $(Atle)$ this is

$$2^{2cn(n-1)+2(a-1)n}\, \cdot\, S_n(a\,, -\frac{1}{2}\,, c) \quad ,$$

times $(-1)^{n+n(n-1)c}$, and I am sure that this annoying extra sign could be explained away. □

References

[CRY] Clara S. Chan, David P. Robbins, and David S. Yuen, *On the volume of a certain polytope*, Experimental Mathematics **9**(2000), 91-99. http://arxiv.org/abs/math/9810154 .

[MeMora] Karola Mészáros and Alejandro H. Morales, *Flow polytopes of signed graphs and the Kostant partition function*, http://arxiv.org/abs/1208.0140, to appear in International Mathematics Research Notices.

[Mora] Alejandro H. Morales, *Open Problems session*, Stanley@70, June 23, 2014, ca. 4:50-5:00pm, http://math.mit.edu/stanley70/Site/Program.html .
Slides are available here: http://math.mit.edu/stanley70/Site/Slides/Morales.pdf

[Morr] William ("Chip") G. Morris, "*Constant Term Identities for Finite and Affine Root Systems: Conjectures and Theorems*", PhD thesis, University of Wisconsin-Madison, 1982. [Advisor: Richard Askey].

[Z] Doron Zeilberger, *Proof of a Conjecture of Chan, Robbins, and Yuen*, Elec. Trans. of Numerical Analysis (ETNA), **9**(1999), 147-148.
http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/cry.html.

Doron Zeilberger, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.
url: http://www.math.rutgers.edu/~zeilberg/ .
Email: zeilberg at math dot rutgers dot edu .

EXCLUSIVELY PUBLISHED IN THE PERSONAL JOURNAL OF SHALOSH B. EKHAD and DORON ZEILBERGER http://www.math.rutgers.edu/~zeilberg/pj.html and arxiv.org .

July 9, 2014